

MatriDerm® – Tissue Regeneration in a One-Step Procedure

Secures the quality of your work

Works with the skills of nature

MatriDerm® is a collagen-elastin scaffold.
Our proprietary Advanced CryoSafe® Technology provides:


- Structural & biochemical properties of native collagen
- No chemical cross-linking – no damage to the matrix
- High similarity to human dermis

MatriDerm® supports all crucial steps for tissue regeneration:

- Immediate cell migration and ingrowth
- Rapid proliferation
- Fast neo-angiogenesis



Scanning-Electron-Microscopy images courtesy of Dr. Mörgelin, Sweden



Angiogenesis 15 days post implantation – courtesy of Prof. Kamolz, Austria

* Split-Thickness Skin Graft

Works for you

Quality in Structure

Native dermal matrix


- Improves functionality through increased skin quality
- Avoids scar formation
- Provides an optimal environment for wound healing:
The reconstructed skin is more elastic and closer to normal skin architecture than STSG* alone

Quality in Efficacy

Permanent wound closure in an easy one-step procedure

- Reduces patient's risk of complications
- Saves time and costs

Effect of MatriDerm® in a Skin Model



Reduced
Wound contraction · Scar formation

Improved
Elasticity · Flexibility · Strength

Works for your patients

Quality of Life

Superior functional and aesthetic outcome

- Restoration of full functionality
- Facilitates early rehabilitation and return to work
- Helps to restore normal daily activity



Photo courtesy of Prof. Kamolz, Austria

MatriDerm® secures the quality of your work

MatriDerm® delivers a superior functional and aesthetic outcome in all kinds of full-thickness wounds – acute and chronic.

3rd degree burn



1 year post-operative



Pressure ulcer



1 week post-operative



Carcinoma excision



6 months post-operative



Mixed chronic ulcer



2 weeks post-operative



Debridement after dog bite 2 years post-operative



Trauma – degloving



6 months post-operative



Adherent scar



1 year post-operative



Trauma – avulsion



5 weeks post-operative



All cases with courtesy of the responsible surgeon